



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

twice I found a strange exception to the usual food of the birds of this family. In two cases I found green olives in the stomach of *Tyrannus verticalis*, and in nearly all birds of the common species *Sayornis nigricans*, killed in winter, I have found pepper berries. *Tyrannus verticalis* is the Bee-martin of California. It flies from its perch near the apiary, captures its prey, then flies back to its perch, making some rapid motion, after which it swallows the bee. I have seen a toad swallow five bees in quick succession; then have killed the toad to find five bee stings sticking to its throat, and as many stingless bees in its stomach. Does the throat of the Kingbird become a sort of pin-cushion for bee stings, or does this bird extract the stings as it manipulates the bee before swallowing? I am eagerly waiting to settle this interesting question. I have taken worker bees from the stomach of the eastern Kingbird (*Tyrannus tyrannus*) but never noticed regarding the disposition of the stings. None of the Kingbirds which I have taken in California have had bees in their stomach, though I am told by good observers that this bird does capture and swallow bees.—A. J. COOK, *Claremont, Cal.*

Rare Visitors to the Connecticut River Valley in Massachusetts in 1895.—*Seiurus motacilla*.—On the 28th day of July, a Louisiana Water Thrush was found dead, on the piazza of a house in the central part of Springfield, undoubtedly having been killed by flying against a window. The capture of a specimen by Dr. J. A. Allen on Mount Tom in April, 1869, is the only other record of its occurrence here.

Rallus elegans.—October 19, a young King Rail was taken in Longmeadow. It was found in the *Zizania aquatica* which grows so profusely along the banks of the Connecticut River. The presence of this species in this part of the valley, I believe, has never before been noted; and the Clapper Rail (*Rallus longirostris crepitans*) has been captured here but twice.

Calcarius lapponicus.—A Lapland Longspur was shot in Longmeadow, November 28.—ROBERT O. MORRIS, *Springfield, Mass.*

A Correction.—In 'The Auk' for April, 1892 (Vol. IX, p. 144), in a note on the 'Habits of the Black-bellied Plover in Massachusetts,' I stated it as my conviction that the *adults* do not assume the gray and white winter plumage after having attained to the full adult spring plumage. I also expressed the same opinion in regard to the plumage of the *adult* Knots in 'The Auk' for January, 1893 (Vol. X, p. 32) in 'Observations on the Knot, *Tringa canutus*.' I now desire to withdraw both of these opinions, as I am inclined to doubt, although not yet *certain*, the correctness of such views, but believe it much better to so state, rather than let a probable error remain to misguide others.—GEORGE H. MACKAY, *Nantucket, Mass.*